

# 2018 Maine Wild Blueberry Pesticide Chart – 2 of 3

**NOTE: THIS CHART IS NOT A SUBSTITUTE FOR READING AND FOLLOWING THE LABEL.** It is unlawful to use any pesticide for other than the registered use. Read and follow the label on the product container. The user assumes all responsibility for use inconsistent with the label. Trade names are used for identification. No product endorsement is implied, nor is discrimination intended against similar materials. Cooperative Extension makes no warranty or guarantee of any kind concerning the use of these products. **Check with your processor regarding PHI restrictions or MRL restrictions for export sales.**



Extremely toxic to fish



Extremely toxic to bees



Extremely toxic to birds



Extremely toxic to people



Moderately toxic to fish



Moderately toxic to bees



Moderately toxic to birds



Moderately toxic to people

## FUNGICIDES

TOXICITY	FUNGICIDE	AI/GROUP	RATE/ACRE	PHI	REI	MAX/ACRE/YR	NOTES
	Equus DF	chlorothalonil <i>Group M5</i>	2.7 to 3.6 lbs	42	12 hrs	10.9 lbs	Apply in prune year for Septoria leafspot or leaf rust at 10-day minimum intervals. Do not add pesticides, fertilizers, or adjuvants unless tested; as phytotoxicity may result. Do not mix with Dipel, Foil, Triton or Latron. High runoff potential for several days to weeks. Human toxicity high for eyes and inhalation.
	Bravo WeatherStik Equus 720 SST		3 to 4 pts			12 pts (9 lbs ai all chlorothalonil)	
	Bumper 41.8 ES Tilt PropiMax EC	propiconazole <i>Group 3</i>	6 oz	30	12 hrs	30 oz (0.84 lb ai all propiconazole products); 2 consecutive applications	Apply at green tip/when >40% stems have blossoms at crown stage for mummy berry. Follow the mummy berry forecast method to time applications or apply every 7 to 10 days following the calendar method. See label for additional disease control. Human toxicity high for skin.
	CaptEvate 68 WDG	captan & fenhexamid <i>Groups M4/17</i>	3.5 to 4.7 lbs	0	48 hrs	21 lbs (3 lbs ai fenhexamid, 35 lbs ai captan)	Apply at 10% bloom for Botrytis. Repeat every 7-10 days through petal fall. Do not make more than 2 consecutive applications. Human toxicity high for skin and eyes.
	Elevate 50 WDG	fenhexamid <i>Group 17</i>	1.5 lbs	0	12 hrs	6 lb (3 lbs ai all fenhexamid products)	Apply at 10% bloom for Botrytis. Apply every 7-10 days when conditions favor disease development, but no more than 2 consecutive applications. See supplemental label.
	Fontelis	penthiopyrad <i>Group 7</i>	16 to 24 oz	0	12 hrs	72 oz per year; 24 oz per application; 2 sequential applications	Apply at green tip/when >40% stems have blossoms at crown stage for mummy berry. Follow the mummy berry forecast method to time applications or apply every 7 to 10 days following the calendar method.
	Indar 2F	fenbuconazole <i>Group 3</i>	6 oz	30	12 hrs	4 applications; 24 oz (0.75 lb ai)	Apply with 1% v/v COC. Apply at green tip/when >40% stems have blossoms at crown stage for mummy berry. Follow the mummy berry forecast method to time applications or apply every 7 to 10 days following the calendar method.
	Luna Tranquility	fluopyram and pyrimethanil <i>Groups 7/9</i>	13.6 to 27 oz	0	12 hrs	54.7 oz per year, 2 sequential applications (0.446 lbs ai all fluopyram products, 2.1 lbs ai all pyrimethanil products)	Apply at green tip/when >40% stems have blossoms at crown stage for mummy berry. Follow the mummy berry forecast method to time applications or apply every 7 to 10 days following the calendar method.

	Pristine  DO NOT TANK MIX – WATER ONLY	pyraclostrobin & boscalid <i>Groups 11/7</i>	18.5 oz	0	12 hrs	4 applications; 92 oz per yr; 23 oz per application* (for all Group 7 or 11 products); 2 sequential applications	Apply at 7-14 day intervals when conditions favor disease development. Avoid application when soil is saturated or when significant rainfall is expected within 48 hrs. Product has high runoff potential for several months. Rotate with a fungicide with a different mode of action. Avoid applying during bloom when bees are present. May only be applied with water.
	Procidic	ascorbic acid, citric acid, lactic acid, glycerin, NaCl N/A	2 to 2.5 oz/10 gal water	0	0 hrs	No maximum	Apply at budbreak and repeat every 7-10 days for <i>Monilinia</i> . Use for powdery mildew only if infection is severe. Spray solution must be pH 5.5-6, low salt concentration, and not chlorinated. Use higher rate during high humidity. Incompatible with dicofol, bipentachlorocyclopentadiene, methomil, cyhexatin, and chloropirites. Do not mix with salt based or high alkali reaction products. Also bactericidal and may be used post-harvest.
	Proline 480 SC	prothioconazole Group 3	5.7 oz	7	12 hrs	2 applications; 11.4 oz; 2 sequential applications	Apply at green tip/when >40% stems have blossoms at crown stage for mummy berry. Follow the mummy berry forecast method to time applications or apply every 7 to 10 days following the calendar method. See label for additional disease control; for <i>Valdensinia</i> suppression only. Product has potential to leach. Human toxicity moderate at longer exposures (see SDS).
 Toxic to shrimp	Quash 50% WDG	metaconazole Group 3	2.5 oz.	7	12 hrs	3 applications, 7.5 oz; 2 sequential applications	Apply at green tip for mummy berry. Follow the mummy berry forecast method to time applications or apply every 7-10 days following the calendar method. Rotate to a product that is not Group 3.
	Quilt Xcel	azoxystrobin & propiconazole <i>Groups 11/3</i>	14 to 21 oz	30	12 hrs	82 oz per yr (0.84 lb ai propiconazole and 0.75 lb ai azoxystrobin); 2 sequential applications	Apply at green tip/40% of flowers buds are at crown stage for mummy berry. Follow the mummy berry forecast method to time applications or apply every 7-10 days following the calendar method. Avoid adjuvants with silicone. Tank-mixing with EC products can lead to phytotoxicity. Rotate to a product that is not Group 11.
	Serenade Max	<i>Bacillus subtilis</i> Not classified	1 to 3 lbs	0	4 hrs	---	For <i>Monilinia</i> suppression, begin application at bud break and repeat every 7-10 days as needed. For Septoria leafspot or leaf rust, begin application prior to disease and repeat every 7-10 days, up to harvest. Spray solution should be pH 4.5-8.5. Rotate with products with other modes of action.
	Switch 62.5 WG	cyprodinil & fludioxonil <i>Groups 9/12</i>	11 to 14 oz	0	12 hrs	56 oz (1.3 lb ai cyprodinil, 0.9 lb ai fludioxonil); 2 consecutive applications	Apply prior to disease for Botrytis and at 7-10 day intervals when conditions favor disease development. Avoid application when soil is saturated or when significant rainfall is expected within 48 hrs, or within 75 ft of water bodies. After 2 consecutive applications, rotate with a fungicide with a different mode of action for 2 applications.

Prepared by Jennifer D'Appollonio, Assistant Scientist; David E. Yarborough, Extension Blueberry Specialist; Seanna Annis, Associate Professor of Mycology; and Rachael Martin, Research Assistant, The University of Maine, Orono, ME 04469. January 2018. **\*Please note: This revised version replaces any previous charts.\***

The University of Maine System is an EEO/AA employer, and does not discriminate on the grounds of race, color, religion, sex, sexual orientation, including transgender status and gender expression, national origin, citizenship status, age, disability, genetic information or veteran's status in employment, education, and all other programs and activities. Please contact the Director of Equal Opportunity, 101 N. Stevens Hall, Orono, ME 04469 at [207-581-1226](tel:207-581-1226) (voice), TTY 711 (Maine Relay System), or [equal.opportunity@maine.edu](mailto:equal.opportunity@maine.edu) with questions or concerns.