

**Always read
the product
label!**

WEED MANAGEMENT SNAPSHOT 2017

UMaine Extension (with help from the UMass Cranberry Chart Book)
Pest Management Office || 491 College Ave || Orono, ME 04473



Herbicides (and active ingredient)	Max. allowable formulation per A	Pre-harvest Interval (PHI)	Restricted Entry Interval (REI)	Max. # of apps	HRAC Group
Callisto® (mesotrione)	16 fl oz	45 days	12 hrs	2	F2
Casoron® 4G (dichlobenil)	100 lb	NA	12 hrs	NA	L
Devrinol® (napropamide)	18 lb or 18 qts	NA	24 hrs	1	K3
Evital® 5G (norflurazon)	160 lb	NA	12 hrs	1	F1
Fusilade® (fluazifop) <i>non-producing beds</i>	N/A	1 year	12 hrs	NA	A
Poast® (sethoxydim)	5 pints	60 days	12 hrs	NA	A
Select Max®, Intensity One® (clethodim)	64 fl oz	30 days	24 hrs	NA	A
Intensity®, Arrow® (clethodim)	32 fl oz	30 days	24 hrs	NA	A
Simazine 4L® (simazine)	2 quarts	NA	12 hrs	1	C1
Reglone® (diquat dibromide) <i>for new bed prep</i>	N/A	1 year	when dry	NA	D
Roundup® (glyphosate)	N/A	30 days	4 hrs	NA	G
Vapam® (metam-sodium) <i>non-producing beds</i>	75 gallons	1 year	5 days	NA	Z

No more Stinger® for Maine cranberries unless we have major problems with clover down the road.

HRAC = Herbicide Resistance Action Committee (rotate amongst the groups to hinder weed resistance)

Before budbreak (~middle of April)

Apply Pre-emergence herbicides prior to vines breaking dormancy. Timing may vary due to weather. **Do not use Callisto® during this period.**

- ✓ Asters, cottongrass, crabgrass, cutgrass, fall panicgrass, ferns, fireweed, haircap moss, loosestrife, mannagrass, marsh St. John's wort, mare's tail (*equisetum*), narrow-leaved goldenrod, needlegrass, nutsedge, pitchfork (beggarstick), ragweed, rattlesnakegrass, rushes, smartweed, sorrel, summergrass, velvetgrass, wild strawberry, woolgrass
- ✓ **[Casoron®]**
- ✓ Pitchforks (beggarsticks), smartweed, ragweed, and summergrass **[Simazine 4L®]**
- ✓ Wild bean/groundnut **[Casoron®, then Devrinol®]**
- ✓ Sphagnum moss **[iron sulfate] [improve drainage]**
- ✓ Barnyardgrass, broadleaf panicgrass, broomsedge, Canada rush, corn grass, cutgrass, dulichium, mare's tail (*equisetum*), needlegrass, nutsedge, smokegrass, spike rush, summergrass, switchgrass, woolgrass **[Evital®]**
- ✓ Brambles / Dewberries **[Late Water, Fall flood, thermal hand-held devices]**
- ✓ Barnyardgrass, cutgrass, fall panicgrass, nutsedge, povertygrass, soft rush, summergrass, warty panicgrass **[Devrinol®]**
- ✓ Dodder **[Casoron®]** or just pull and flag if small areas; Callisto® later in the season

~May to 45 days before harvest or flooding *(but not during bloom)*

- ✓ Amaranth, atriplex, buffalobur, carpetweed, carrot, chickweed, cocklebur, crabgrass, dodder, horse nettle, jimsonweed, lambsquarters, mustard, nightshade, pigweed, ragweed, smartweed. **When applied early:** some annual and perennial grasses (*but probably better results if followed up with a grass herbicide*), most annual broadleaves, many herbaceous perennial broadleaves such as yellow loosestrife*, silverleaf, asters, rushes, birdsfoot trefoil, St. john's wort, and sedges, and some woody perennials such as blackberry. **[Callisto®] (max of 2 applications per season, whether or not the full rate is used either time) (Permanent control of many perennial weeds may take several years of Callisto® treatment but avoid using it every year if possible)** *Level of yellow loosestrife and dodder control has been variable in Massachusetts. Resistance to Callisto® has been reported in other crops after 7 years of use. Therefore, avoid using Callisto® year after year!

-
- June through August**
- ✓ **many broadleaved weeds [Callisto®] (45-day PHI)**
 - ✓ Cinnamon fern, feather fern, sensitive fern [**iron sulfate and salt**]
 - ✓ Asters, cinquefoil, feather fern, sensitive fern, marsh St. John's wort [**iron sulfate**]
 - ✓ Actively growing true grasses (annual & perennial) on beds [**Poast®**] [**Select Max®**]
 - ✓ Ditch weeds (in dry ditches) [**Roundup®** or **Rodeo®**] [**Flame cultivation, mowing, or pulling by hand early in the spring & early summer**]; **duckweed** [**draining the ditch**]
 - ✓ Tall weeds on beds [**Roundup®**]
 - ✓ Wild bean / groundnut [**salt solution**]
 - ✓ Dodder and poison ivy [**spot treatments with concentrated Callisto®**]
-

- Weeds outside of the beds**
- ✓ **Summer:** Water weeds in farm ponds and reservoirs such as bladderwort, coontail, elodea, and pondweeds [**Diquat®**] [**Reward®**] Water temperature should be over 50°F for best activity. Do not use in or on bog ditches!
 - ✓ **Summer:** Effective choice for cattails and water lilies [**Rodeo® + nonionic surfactant**] (**but cannot use in interior ditches**)
 - ✓ Woody and broadleaf perennial weeds (**non-bed use only!!**) [**Weedone CB®**] Weedone is no longer produced, but available product may be used; apply in March, if possible; spray to wet and avoid drift onto the bed! [**Crossbow®**] Crossbow contains trichlopyr for which there is **no food tolerance**, so be careful using this product – use it only on weeds located far away from the bed(s) because it has the potential to drift (nearby plants such as apple trees, grapes, tomatoes, etc. may also be injured from any drifting)
-

- In the Fall After Harvest**
- ✓ Aster, blue joint, cottongrass, Carex spp. (“true” sedges), cutgrass, loosestrife, marsh St. John's wort, mud rush, narrow-leaved goldenrod, needlegrass, nutsedge, ragweed, sphagnum moss, summergrass, woolgrass [**Casoron®**]
 - ✓ Barnyardgrass, Canada rush, fall panicgrass, needlegrass, nutsedge, summergrass, povertygrass (plus winter sanding), and switchgrass [**Evital®**]
 - ✓ Barnyardgrass, cutgrass, fall panicgrass, nutsedge, povertygrass, spike rush [**Devrinol®**]
 - ✓ **Spot-treatments [such as with Roundup WeatherMAX®, Evital®]**
 - ✓ Pitchforks, smartweed (ladythumb), ragweeds, & summergrass [**Simazine 4L®**]
-

- Bed Renovation or New Bed Preparation (non-producing beds only)**
- Soil fumigants: **Basamid®** (dazomet), **Vapam®** (metam-sodium) **Note:** Both of these products are in the same HRAC group → Group Z.
 - **Reglone®** (diquat dibromide) + **nonionic surfactant** (intended for site preparation; cannot be applied by chemigation; use 1.5-2 pints/15 gallons of water by ground; may need repeat applications)
 - **Fusilade®** (fluazifop) post-emergence; for true grasses on non-producing beds; Use 0.75 fl oz. per gallon of water; grasses will turn yellow and die back over a 1 to 4-week period depending on climatic conditions; add a COC at 0.5-1% v:v or 0.25-0.5% v:v if using a NIS.

A Member of the University of Maine System